

BREAKOUT

The Newsletter of the Hastings and
Napier Amateur Radio Clubs

Hastings Branch 13 NZART – Napier Branch 25 NZART

Volume 8, Issue 6, June 2010



Hastings Br
13
Club Calls
ZL2AS
ZL2QS

Napier Br 25
Club Call
ZL2GT

IRLP
Node
6793
147.250

Branch
Nets
9.00 AM
Sunday
Morning
3615 Hz
147.250
MHz

Editor
John Newson
ZL2VAF



ZL2LF caught this Harvard doing unusual aerobatics at the recent Napier Air Show. "I was extremely lucky to capture this moment, it was so short that very few people actually saw it. BUT- with the wonder of modern technology, I managed it."
Ed:- I am not sure the fee paid to the pilot but I am sure if you approached ZL2LF he would be able to arrange your callsign displayed at a future air show.

<http://groups.yahoo.com/group/zl2as/>



Join the KIWI DX Group
Talk to ZL2AL for Details

Inside This Issue

Hastings Branch 13 Report	Page 2
Napier Branch 25 Report	Page 3
Internet Link – Yaesu Manuals	Page 3
Amateur Operator Charged	Page 4
Whats Wrong With The Sun?	Page 7
Notices	Page 8
Buy, Sell or Exchange	Page 8

HASTINGS BRANCH 13

President: Warren Harris ZL2AJ 027 564 9284 or 929 9088 email warren@technaserve.com
Vice President: Robert Wallace ZL2SG Ph. 878 4993 email ffonzrjw@xnet.co.nz
Secretary: David Walker ZL2DW Ph 8760518, email david@apexradiocomms.co.nz
Treasurer: Bill Lowes ZL2UBG Ph. 877 5078 email bill.lowes@xnet.co.nz
AREC/CD: Robert Wallace ZL2SG Ph. 878 4993 email ffonzrjw@xnet.co.nz
AREC Deputy: Warren Harris ZL2AJ 027 564 9284 or 929 9088 email warren@technaserve.com

Committee: - Lee Jennings ZL2AL Ph. 844 1226 email leejen@paradise.net.nz
- Peter Dingley ZL2LF Ph. 843 2664 email peter.dingley@paradise.net.nz
- Charlotte Shuker ZL2QC Ph. 929 9088
- Rob Leicester ZL2 RFL ph 8786381 wk 8782828 email zl2rfl@yahoo.co.nz

Hastings QSL Distribution: Chris Johnson ZL2VC Ph.879 5219 email zl2vc@xtra.co.nz
Magazine Editor: John Newson ZL2VAF Ph. 027 230 3642 email john@thecomputerman.co.nz
NZART License Examiners: Lee - ZL2AL and Peter - ZL2LF
Club Call: ZL2AS and ZL2QS

Club Nights: Fourth Wednesday each month at 7.30 pm Surf Club Rooms, Windsor Park, Hastings

Hastings Branch 13 - President's Report

Hi all.

This Wednesdays meeting is an away meeting. It will be at the Faraday Centre, Faraday St Napier. Please meet there at 1930 hours. Bring \$6 and a jersey - there is no heating in the exhibit hall. It should be interesting and exciting for all. Bring the xyl.

Call for operators. The Gisborne Rally is on the 7th of August. I am looking for operators who would like to provide comms for this event. Please contact me directly for details.

NZART Conference. I attended this on Br13's and Br25' behalf, and proxy for Br60 (Taupo). The remits were voted on and the outcomes were as follows.

- Remit 1 (ARX free) lost on vote count
- Remit 2 (Family members) Passed on voice
- Remit 3 (XYZ calls) Passed on voice
- Remit 4 (Vote for ELL) Passed on voice
- Remit 5 (AREC forum time) Lost on voice
- Remit 6 (Constitution to effect remit 2) assed on voice

Entry Level Licence mandate (as per remit 4) Lost on vote count. A foundation licence is now off the table.

Other issues from the weekend. There was a motion on the floor that we seek to raise our legal power limit. In light of the recent prosecution of a ham for breaching the existing rules it was thought that now was not the right time. Lost on voice.

Maybe a remit next year?

During the weekend I saw presentations on SDR radio, Digital modes software and VHF oscillators. I attended forums on IRLP and APRS, as well as the open forum. The idea of our expo contest? session gained some traction and interest and I will be pushing this to happen.

I have also been formally asked to head a committee and be the national training officer. I have accepted this position and with John Andrews, Vaughan Henderson and Tony Case, we will be formulating plans and resources to assist branches to train new amateur radio operators.

Next years conference will be at Upper Hutt.

See you at the Faraday Centre on Wednesday.

Regards,

Warren Harris ZL2AJ

NAPIER BRANCH 25

President: Laurie Winton ZL2TC 843 8519

Secretary: Stan White ZL2ST 845 2422

Treasurer: Stan White ZL2ST 845 2422

AREC: Tony Wall ZL2RZ

email laurie@wilcom.co.nz

email stan.white@clear.net.nz

email stan.white@clear.net.nz

email ttwall@paradise.net.nz

Committee:

Tony Wall ZL2RZ 8354424

Gary James ZL2GAZ 843 9596

Michael ZL2FAR 843 4210

email ttwall@paradise.net.nz

email gazzaj@paradise.net.nz

Committee Meetings: Third Monday of the month 7pm at Club Rooms

Club Call: **ZL2GT**

Club Nights: First Wednesday each month (except January) 7.30pm at the Club Rooms:
123 Latham Street Napier

NAPIER NEWS...

The next Napier meeting will be on Wednesday 7 July at the clubrooms at 7.30 pm. We will have a digital modes demonstration in preparation for a kitset construction evening later in the month. Lee has acquired all of the parts for the kitset project which is a computer – rig interface for digital modes. The manufacture of a PC board under Lee's direction will be part of the construction evening. If you are interested in building a kitset come along to the July meeting. Hastings members welcome.

Thanks to Warren ZL2AJ for representing us at conference at Auckland.

HF band conditions are still poor with the only QSO this month being with a YN in Nicaragua on 30M CW. Last month provided me with a new country – ET3SID Sid in Addis Ababa, Ethiopia on 20M. I have been looking for ET for a number of years and finally clicked at around 4 pm about 2 months ago when he was looking for ZLs. There are only 3 or 4 hams there who operate spasmodically.

Stan ZL2ST



Internet Link

Yaesu Service Manuals free download

<http://www.kb3lsr.com/yaesu.htm>

Amateur radio operator charged

Radio Spectrum Management recently prosecuted Amateur Radio Operator Alan Potter for transmitting outside the terms and conditions of the General User Radio Licence for Amateur Radio Operators.

The radio spectrum is an important resource for New Zealand and a key part of Radio Spectrum Management's work is to ensure responsible use of the radio spectrum in a way that doesn't cause interference to other users.

All amateur radio operators are required to gain a qualification that covers the risks and harmful effects of interference from their transmitting equipment. This includes the risks associated with transmitting at high power levels.

"As a qualified amateur operator, Mr Potter is well aware of the risks but has chosen to ignore them", says Chris Brennan, Compliance Manager for Radio Spectrum Management.

"This behaviour is unacceptable. Radio Spectrum Management are serious about protecting the radio spectrum for all users; we are continually monitoring and enforcing radio spectrum compliance, which includes prosecution when necessary". Radio Spectrum Management was alerted to a video Mr Potter posted on 'You Tube', a well known online video sharing site, which showed his transmitter operating at 3100 Watts. Operating at such high transmitting power is likely to cause interference to, and disruption of, a range of other licensed radio services in the local area.

Radio Spectrum Management's role is to protect the public good by ensuring equipment capable of transmitting radio waves complies with the terms, conditions and restrictions of radio licences and International standards applicable in New Zealand.

Conviction and charge

Mr Potter was charged in the Christchurch District Court on 14 April. He was found guilty of breaching section 113 of the Radiocommunications Act 1989. He was fined \$1,750 and \$130 for costs, plus he has been required to forfeit his radio equipment.

For the purposes of section 113, any person who erects, constructs, establishes, maintains, or is in possession of any radio transmitter is presumed to have used the radio transmitter. In this case, Mr Potter was found in possession of radio transmitting equipment that was capable of operating at a significantly higher power than the Amateur Radio Operators General Licence allows.

What's wrong with the sun?

SUNSPOTS come and go, but recently they have mostly gone. For centuries, astronomers have recorded when these dark blemishes on the solar surface emerge, only for them to fade away again after a few days, weeks or months. Thanks to their efforts, we know that sunspot numbers ebb and flow in cycles lasting about 11 years.

But for the past two years, the sunspots have mostly been missing. Their absence, the most prolonged for nearly a hundred years, has taken even seasoned sun watchers by surprise. "This is solar behaviour we haven't seen in living memory," says David Hathaway, a physicist at NASA's Marshall Space Flight Center in Huntsville, Alabama.

The sun is under scrutiny as never before thanks to an armada of space telescopes. The results they beam back are portraying our nearest star, and its influence on Earth, in a new light. Sunspots and other clues indicate that the sun's magnetic activity is diminishing, and that the sun may even be shrinking. Together the results hint that something profound is happening inside the sun. The big question is what?

The stakes have never been higher. Groups of sunspots forewarn of gigantic solar storms that can unleash a billion times more energy than an atomic bomb. Fears that these [giant solar eruptions could create havoc on Earth](#), and disputes over the sun's role in climate change, are adding urgency to these studies. When NASA and the European Space Agency launched the [Solar and Heliospheric Observatory](#) almost 15 years ago, "[understanding the solar cycle](#) was not one of its scientific objectives", says Bernhard Fleck, the mission's project scientist. "Now it is one of the key questions."

- Sun behaving badly

Sunspots are windows into the sun's magnetic soul. They form where giant loops of magnetism, generated deep inside the sun, well up and burst through the surface, leading to a localised drop in temperature which we see as a dark patch. Any changes in sunspot numbers reflect changes inside the sun. "During this transition, the sun is giving us a real glimpse into its interior," says Hathaway.

When sunspot numbers drop at the end of each 11-year cycle, solar storms die down and all becomes much calmer. This "solar minimum" doesn't last long. Within a year, the spots and storms begin to build towards a new crescendo, the next solar maximum.

What's special about this latest dip is that the sun is having trouble starting the next solar cycle. The sun began to calm down in late 2007, so no one expected many sunspots in 2008. But computer models predicted that when the spots did return, they would do so in force. Hathaway was reported as thinking the next solar cycle would be a "doozy": more sunspots, more solar storms and more energy blasted into space. Others [predicted](#) that it would be the most active solar cycle on record. The trouble was, no one told the sun.

The first sign that the prediction was wrong came when 2008 turned out to be even calmer than expected. That year, the sun was spot-free 73 per cent of the time, an extreme dip even for a solar minimum. Only the minimum of 1913 was more pronounced, with 85 per cent of that year clear.

As 2009 arrived, solar physicists looked for some action. They didn't get it. The sun continued to languish until mid-December, when the largest group of sunspots to emerge for several years appeared. Finally, a return to normal? Not really.

Even with the solar cycle finally under way again, the number of sunspots has so far been well below expectations. Something appears to have changed inside the sun, something the models did not predict. But what?

The flood of observations from space and ground-based telescopes suggests that the answer lies in the behaviour of two vast conveyor belts of gas that endlessly cycle material and magnetism through the sun's interior and out across the surface. On average it takes 40 years for the conveyor belts to complete a circuit (see diagram).

When Hathaway's team looked over the observations to find out where their models had gone wrong, they noticed that the conveyor-belt flows of gas across the sun's surface have been speeding up since 2004.

The circulation deep within the sun tells a different story. Rachel Howe and Frank Hill of the National Solar Observatory in Tucson, Arizona, have used observations of surface disturbances, caused by the solar equivalent of seismic waves, to infer what conditions are like within the sun. Analysing data from 2009, they found that while the surface flows had sped up, the internal ones had slowed to a crawl.

These findings have thrown our best computer models of the sun into disarray. "It is certainly challenging our theories," says Hathaway, "but that's kinda nice."

It is not just our understanding of the sun that stands to benefit from this work. The extent to which changes in the sun's activity can affect our climate is of paramount concern. It is also highly controversial. There are those who seek to prove that the solar variability is the major cause of climate change, an idea that would let humans and their greenhouse gases off the hook. Others are equally evangelical in their assertions that the sun plays only a minuscule role in climate change.

If this dispute could be resolved by an experiment, the obvious strategy would be to see what happens when you switch off one potential cause of climate change and leave the other alone. The extended collapse in solar activity these past two years may be precisely the right sort of test, in that it has significantly changed the amount of solar radiation bombarding our planet. "As a natural experiment, this is the very best thing to happen," says Joanna Haigh, a climatologist at Imperial College London. "Now we have to see how the Earth responds."

- The climate link

Mike Lockwood at the University of Reading, UK, may already have identified one response - the unusually frigid European winter of 2009/10. He has [studied records covering data stretching back to 1650](#), and found that severe European winters are much more likely during periods of low solar activity ([New Scientist, 17 April, p 6](#)). This fits an emerging picture of solar activity giving rise to a small change in the global climate overall, yet large regional effects.

Another example is the Maunder minimum, the period from 1645 to 1715 during which sunspots virtually disappeared and solar activity plummeted. If a similar spell of solar inactivity were to begin now and continue until 2100, it would [mitigate any temperature rise through global warming by 0.3 °C on average](#), according to calculations by [Georg Feulner](#) and [Stefan Rahmstorf](#) of the Potsdam Institute for Climate Impact Research in Germany. However, something amplified the impact of the Maunder minimum on northern Europe, ushering in a period known as the Little Ice Age, when colder than average winters became more prevalent and the average temperature in Europe appeared to drop by between 1 and 2 °C.

A corresponding boost appears to be associated with peaks in solar output. In 2008, Judith Lean of the Naval Research Laboratory in Washington DC published a study showing that high solar activity has a disproportionate warming influence on northern Europe ([Geophysical Research Letters, vol 35, p L18701](#)).

So why does solar activity have these effects? Modellers may already be onto the answer. Since 2003, spaceborne instruments have been measuring the intensity of the sun's output at various wavelengths and looking for correlations with solar activity. The results point to the sun's emissions of ultraviolet light. "The ultraviolet is varying much, much, much more than we expected," says Lockwood.

Ultraviolet light is strongly linked to solar activity: solar flares shine brightly in the ultraviolet, and it helps carry the explosive energy of the flares away into space. It could be particularly significant for the Earth's climate as ultraviolet light is absorbed by the ozone layer in the stratosphere, the region of atmosphere that sits directly above the weather-bearing troposphere.

More ultraviolet light reaching the stratosphere means more ozone is formed. And more ozone leads to the stratosphere absorbing more ultraviolet light. So in times of heightened solar activity, the stratosphere heats up and this influences the winds in that layer. "The heat input into the stratosphere is much more variable than we thought," says Lockwood.

Enhanced heating of the stratosphere could be behind the heightened effects felt by Europe of changes in solar activity. Back in 1996, Haigh showed that the temperature of the stratosphere influences the passage of the jet stream, the high-altitude river of air passing from west to east across Europe.

Lockwood's latest study shows that when solar activity is low, the jet stream becomes liable to break up into giant meanders that block warm westerly winds from reaching Europe, allowing Arctic winds from Siberia to dominate Europe's weather.

The lesson for climate research is clear. "There are so many weather stations in Europe that, if we are not careful, these solar effects could influence our global averages," says Lockwood. In other words, our understanding of global climate change could be skewed by not taking into account solar effects on European weather.

Just as one mystery begins to clear, another beckons. Since its launch 15 years ago, the SOHO spacecraft has watched two solar minimums, one complete solar cycle, and parts of two other cycles - the one that ended in 1996 and the one that is just stirring. For all that time its [VIRGO instrument](#) has been measuring the total solar irradiance (TSI), the energy emitted by the sun. Its measurements can be stitched together with results from earlier missions to provide a 30-year record of the sun's energy output. What this shows is that during the latest solar minimum, the sun's output was 0.015 per cent lower than during the previous lull. It might not sound like much, but it is a hugely significant result.

We used to think that the sun's output was unwavering. That view began to change following the launch in 1980 of NASA's [Solar Maximum Mission](#). Its observations show that the amount of energy the sun puts out varies by around 0.1 per cent over a period of days or weeks over a solar cycle.

- **Shrinking star**

Despite this variation, the TSI has dipped to the same level during the three previous solar minima. Not so during this recent elongated minimum. Although the observed drop is small, the fact that it has happened at all is unprecedented. "This is the first time we have measured a long-term trend in the total solar irradiance," says Claus Fröhlich of the [World Radiation Centre](#) in Davos, Switzerland, and lead investigator for the VIRGO instrument.

If the sun's energy output is changing, then its temperature must be fluctuating too. While solar flares can heat up the gas at the surface, changes in the sun's core would have a more important influence on temperature, though calculations show it can take hundreds of thousands of years for the effects to percolate out to the surface. Whatever the mechanism, the cooler the surface, the less energy there is to "puff up" the sun. The upshot of any dip in the sun's output is that the sun should also be shrinking.

Observations suggest that it is - though we needn't fear a catastrophe like that depicted in the movie *Sunshine* just yet. Back in the 17th century French astronomer Jean Picard made his mark by measuring the sun's diameter. His observations were carried out during the Maunder minimum, and he obtained a result larger than modern measurements. Was this simply because of an error on Picard's part, or could the sun genuinely have shrunk since then? "There has been a lot of animated discussion, and the problem is not yet solved," says Gérard Thuillier of the Pierre and Marie Curie University in Paris, France.

Observations with ground-based telescopes are not precise enough to resolve the question, due to the distorting effect of Earth's atmosphere. So the French space agency has designed a mission, aptly named [Picard](#), to return precise measurements of the sun's diameter and look for changes.

Frustratingly the launch, on a Russian Dnepr rocket, is mired in a political disagreement between Russia and neighbouring Kazakhstan. Until the dispute is resolved, the spacecraft must wait. Every day of delay means valuable data being missed as the sun takes steps, however faltering, into the next cycle of activity. "We need to launch now," says Thuillier.

What the sun will do next is beyond our ability to predict. Most astronomers think that the solar cycle will proceed, but at significantly depressed levels of activity similar to those last seen in the 19th century. However, there is also evidence that the sun is inexorably losing its ability to produce sunspots ([see "The sunspot forecast"](#)). By 2015, they could be gone altogether, plunging us into a new Maunder minimum - and perhaps a new Little Ice Age.

Of course, solar activity is just one natural source of climate variability. Volcanic eruptions are another, spewing gas and dust into the atmosphere. Nevertheless, it remains crucial to understand the precise changeability of the sun, and the way it influences the various regional patterns of weather on Earth. Climate scientists will then be able to correct for these effects, not just in interpreting modern measurements but also when attempting to reconstruct the climate stretching back centuries. It is only by doing so that we can reach an unassailable consensus about the sun's true level of influence on the Earth and its climate.

NOTICES

Gisborne Car Rally
7th August



Hawkes Bay Car Rally
28th August



Rally Wairarapa
11th & 12th September



SAREX
2nd & 3rd October



*Please feel free to send notices to
john@thecomputerman.co.nz*

Buy – Sell - Etc

*Please feel free to send you Buy, Sell or Trade
notices to john@thecomputerman.co.nz
Free listing to Branch Members*